



Vexoderm
Dressing

CLINICAL EVIDENCE VEXODERM DRESSING

Skin and wound care products with
antioxidant and regenerative properties.

www.vexoderm.com

CLINICAL EVIDENCE

1st Vexoderm

VERY DEEP EROSIVE LESIONS OF TRAUMATIC ORIGIN

Clinical Cases Contest

Tarta

Turtle. 10 years old.

Previous treatment

No previous pathologies.

Dr.ssa Stefania Di Paco, DVM,

Dr.ssa Raffaella Gambirasio, DVM &

Dr.ssa Marzia Possenti, DVM

Treatment for 6 weeks with **Vexoderm**

Dressing and **Vexoderm Gel+**

BEGINNING



Both forelimbs had very deep and contaminated erosive lesions, with significant soft tissue loss and consequent exposure of forearm bone structures.

WEEK 2



For the forelimbs, every 7 days the application of **Vexoderm Dressing**, is covered with greasy gauze, cotton wool, and a self-adhesive bandage.

For the tail, it was decided to use **Vexoderm Gel+**



WEEK 6



The gradual and steady reconstruction of the different tissue layers achieved excellent wound healing.

Complete healing occurred with the application of **Vexoderm Gel+** on the forelimb wounds as well.



artinvet



vexoderm

CLINICAL EVIDENCE

LACERATED WOUND DUE TO TRAFFIC ACCIDENT

1st Vexoderm

Clinical Cases Contest

Gilda

Cat. 6 years old.

Previous treatment

No previous pathologies.

Dott. Alessandro Negro, DVM

Vexoderm Dressing is kept in place for 7 days. Treatment continues with the application of **Vexoderm Gel+** every 8 hours by the owner.

BEGINNING



Chronic wound contaminated by rubbing. Surgical debridement of the lesion is performed.



WEEK 0



Application of **Vexoderm Dressing**. Kept in place for 7 days.

WEEK 1



Application of **Vexoderm Gel+** every 8 hours by the owner.

WEEK 2



Contraction of the hock wound, which appears drier.

WEEK 3



Complete healing. Application of **Vexoderm Gel+** every 8 hours by the owner continues to improve scar healing.



artinvet



vexoderm

CLINICAL EVIDENCE

LACERATED-BRUISED WOUND FROM RUBBING

Colony cat

Female. Adult.

Previous treatment:

No previous pathology

Dott. Alessandro Negro, DVM, MSc

Treatment takes 4 weeks, combining

Vexoderm Dressing and **Vexoderm Gel+**.

The use of the **Vexoderm Gel+** starts at week 3. Soft tissue surgery and wound care. Medical and surgical oncology.

BEGINNING



Application of **Vexoderm Dressing**

WEEK 1



Cleaning of the wound bed and replacement of the dressing. Very lively granulation. Margins begin to contract.

WEEK 1,5



Very evident wound contraction. The volunteer proposes to apply **Vexoderm Gel+** every 8 hours to avoid further sedation.

WEEK 2



Tissues are closed to healing. Without sedation, getting too close is risky.

WEEK 3



The wound is completely healed.



artinet



vexoderm

CLINICAL EVIDENCE

EXPLODED ABSCESS IN THE SACRO-ISCHIAL REGION

Golgo

European Shorthair cat, sterilised female. 9 years old.

Dott. Alessandro Negro, DVM, MSc

Soft tissue surgery and wound care.
Medical and surgical oncology.

Previous treatment:

Attempt to treat it with sugar.
Sedation, trichotomy, washing and cleansing. Disinfection with chlorhexidine and povidone-iodine.
Surgical debridement and buttonholes for tie-overs.

Application of **Vexoderm Dressing** and tie-over. Excellent granulation and contraction of margins.

BEGINNING



WEEK 1



WEEK 2



WEEK 2



CLINICAL EVIDENCE

TISSUE LOSS DUE TO TRAUMATISM

Dog

Female. 3 months old.

Previous treatment

Manuka honey plus antimicrobial cream

Vexoderm Dressing and Vexoderm Gel+ treatment

Vexoderm Dressing 10x7, 9x4


VetCenter Clinic, Bucharest.

Dr. Isabelle Caloian.


BEGINNING



The patient is sedated, the wound is cleansed, and a proximal suture is applied with resorbable thread.



A small amount of **Vexoderm Gel+** is applied using a cotton swab.



The client applies **Vexoderm Gel+** at home morning and evening using a cotton swab.

WEEK 1



The client continues to apply **Vexoderm Gel+** twice a day.



The feathers are starting to grow back. Keeps applying **Vexoderm Gel+** until complete healing.

WEEK 2



Healing is completed. The head feathers have completely grown.



artinvet



vexoderm

CLINICAL EVIDENCE

EXPOSED BONE DUE TO TRAUMATISM

European Shorthair cat

Sterilised female. 9 years old.

Previous treatment:

None

Vexoderm Dressing treatment

No. of changes: 4 (once a week).

No. of dressings: 4 Vexoderm Dressing 9x4.

VetCenter Clinic, Bucharest.

Dr. Isabelle Caloian.

BEGINNING



WEEK 1



WEEK 2



WEEK 3



CLINICAL EVIDENCE

TRAUMATIC WOUND IN ONE LEG

Fox Terrier

Dog, male. 10 years old.

Previous treatment:

No previous pathology

Vexoderm Dressing treatment

No. of changes: 3 (once a week).

No. of dressings: 3 Vexoderm 9x4.

Wound closure: week 3.

BEGINNING



WEEK 1



WEEK 2



WEEK 3



artinvet



vexoderm

CLINICAL EVIDENCE

TRAUMATIC WOUND IN ONE LEG

Border Collie

Female. 11 months old.

Previous treatment

None

Vexoderm Dressing treatment

No. of changes: 3 (once a week).

No. of dressings: 3 Vexoderm 9x4.

Wound closure: week 3.


BEGINNING



WEEK 1



WEEK 2



WEEK 4



WEEK 5



WEEK 6



CLINICAL EVIDENCE

SURGICAL RESECTION OF HEPATOID GLAND TUMOUR IN THE TAIL

Golden Retriever

Dog, male. 9 years old.

Previous treatment:

Antibiotics

Vexoderm Dressing treatment

No. of changes: 2 (once a week).

No. of dressings: 2 Vexoderm 9x4.

Wound closure: week 2.

Prevents potential wound dehiscence.

No maceration

BEGINNING



WEEK 1



WEEK 2



WEEK 10



artinvet



vexoderm

CLINICAL EVIDENCE

SUPPURATIVE NECROTIC ULCER. INFECTED.

American Staffordshire

Dog, female. 2 years old.

Previous treatment

Antibiotics / NSAIDs. Surgical debridement.

Vexoderm Dressing treatment

No. of changes: 7 (first change after 3 days and then once a week).

No. of dressings: 7 Vexoderm 9x4.

Wound closure: week 7.

BEGINNING



WEEK 1



WEEK 2



WEEK 4



WEEK 7



6 MONTHS



CLINICAL EVIDENCE

BURN CAUSED BY DELIVERATELY PUTTING THE CAT IN THE OVEN.

Felis Vulgaris

Cat, female. 10 months old.

Previous treatment

Moisturising products. Antibiotics.

Surgical debridement.

Vexoderm Dressing treatment

No. of changes: 9 (two changes the first week and then once a week).

No. of dressings: 12 Vexoderm 10x7.

2 Vexoderm 9x4.

Wound closure: week 8.


BEFORE DEBRIDAMENT



BEGINNING




WEEK 1



WEEK 2



WEEK 3



WEEK 4



WEEK 8



WEEK 8



CLINICAL EVIDENCE

TISSUE LOSS DUE TO TRAUMA

European Shorthair

Cat, female. 1 year old.

Trauma 10 days in evolution

Vexoderm Dressing treatment

No. of changes: 5 (once a week).

No. of dressings: 5 Vexoderm 9x4.

Wound closure: week 5.


Previous treatment

Antibiotics. Saline cures

BEGINNING



WEEK 1



WEEK 1



WEEK 2



WEEK 3




WEEK 4



WEEK 5



WEEK 6



artinvet



vexoderm

CLINICAL EVIDENCE

POLYTRAUMATISM, NECROTIC WOUND, EMPHYSEMA AND EDEMA

Cat

Male. 7 years old.

Previous treatment

Suture / Spray (silver)

Vexoderm Dressing treatment

No. of changes: 10 (once a week).

No. of dressings: 10 Vexoderm 9x4.

Wound closure: week 10.

BEGINNING



WEEK 1



WEEK 2



WEEK 3



WEEK 5




WEEK 9



WEEK 9



WEEK 10



CLINICAL EVIDENCE

CUT, HYPERGRANULATION “PROUD FLESH”

Crossbreed horse

Male. 15 years old.

Previous treatment

Antibiotics and cures. Surgical debridement.

Vexoderm Dressing treatment

No. of changes: 32 (once a week).

No. of dressings: 24 Vexoderm 10x7 and 16 Vexoderm 9x4.

Wound closure: week 32.

Tissue resection at weeks -1, 2, 4 & 5

BEGINNING



WEEK 1



WEEK 4



WEEK 10



WEEK 17



WEEK 18



6 months later



CLINICAL EVIDENCE

CHRONIC WOUND

Thoroughbred horse

Male. 2 years old. 5 months in evolution.

Previous treatment

Antibiotics 2 days before Vexoderm

Vexoderm Dressing treatment

No. of changes: 5 (every 7-10 days).

No. of dressings: 5 Vexoderm 10x7.

Wound closure: week 7.

BEGINNING



WEEK 2



WEEK 3



WEEK 4




WEEK 6



WEEK 7



1 month later



CLINICAL EVIDENCE

CHRONIC ULCER OF UNKNOWN ETHIOLOGY.

Horse

Previous treatment

Multiple different treatments including honey.

Vexoderm Dressing treatment

No. of changes: 7 (once a week).

No. of dressings: 4 Vexoderm 10x7.

6 Vexoderm 9x4.

Wound closure: treatment with Vexoderm ended before wound closure (week 7).

BEGINNING



WEEK 1



WEEK 2



WEEK 3



WEEK 4



WEEK 6



WEEK 7



CLINICAL EVIDENCE

CUT WITH FLAP

Thoroughbred horse

Female. 22 years old. 5 months in evolution.

Previous treatment

Antibiotics and anti-inflammatory drugs.

Vexoderm Dressing treatment

No. of changes: 7 (once a week).

No. of dressings: 7 Vexoderm 10x7.

Wound closure: week 7.

FLAP REMOVAL



BEGINNING



WEEK 2



WEEK 3



WEEK 5



WEEK 7



CLINICAL EVIDENCE

ULCER DUE TO LAPAROTOMY. POSSIBLE REACTION TO SUTURE

Bulldog

Dog, female. 4 years old.

Previous treatment:


Systemic antibiotics. Dressing 6 days after surgery

Vexoderm Dressing treatment

Wound activation and healing.

Treatment with Vexoderm Dressing until wound closure.

BEGINNING



WEEK 2



WEEK 3



CLINICAL EVIDENCE

CHRONIC SADDLE SORE IN BOTH SIDES FOR OVER 5 MONTHS.

Quarter Horse

Horse, male. 12 years old.

Previous treatment:

Ointment

Vexoderm Dressing treatment

Vexoderm Dressing is applied with dressing changes once a week for two weeks until the wounds are completely closed. A few weeks later hair growth is observed.

BEGINNING



WEEK 2



WEEK 3

